

ECOLOGY OF ORAL BIO-FILM

Dr. Tirthankar Debnath*, Dr. Pradip Kumar Giri**

ABSTRACT

Most microbial ecosystems contain large numbers of genetically distinct microorganisms, and the human mouth is not different. The biofilm communities are complex and dynamic structures that accumulate through the sequential and ordered colonization of multiple oral bacteria. Oral biofilms are characterized by surface attachment, structural heterogeneity, complex interspecies interactions, and an extracellular matrix of polymeric substances, and are high-density micro-niches that differ dramatically from surrounding conditions. Here, it is an attempt to discuss the ecology of oral biofilm.

KEY WORDS

Microbial ecosystem, biofilm, quorum sensing

ABOUT THE AUTHORS

*Assistant Professor, **Associate Professor
Department of Periodontology
Dr. R. Ahmed Dental College & Hospital, Kolkata

CORRESPONDING AUTHOR

Dr. Pradip Kumar Giri
Associate Professor, Department of Periodontology
Dr. R. Ahmed Dental College & Hospital, Kolkata

INTRODUCTION

The biofilm communities are complex and dynamic structures that accumulate through the sequential and ordered colonization of multiple oral bacteria (Kolenbrander et al., 2002). One of the most notable features of oral biofilms is that oral bacteria growing in the biofilms frequently express phenotypes that are different from those of planktonic bacteria. Oral biofilms are characterized by surface attachment, structural heterogeneity, complex interspecies interactions, and an extracellular matrix of polymeric substances, and are high-density micro-niches that differ dramatically from surrounding conditions. Bacterial species present in the oral biofilm communities interact cooperatively or competitively with other members. It has been shown that the bacterial interactions that influence biofilm formation, metabolic change and physiological function involve various different mechanisms (Fig:1). From a physical aspect, planktonic bacterial cells attach directly to surfaces of the oral cavity or bind indirectly to other bacterial cells that have already colonized (Kolenbrander et al., 2002). From metabolic and physiological points of view, avenues of communication within oral biofilms are likely to include metabolic communication, genetic exchange, quorum-sensing, etc. (Chalmers et al., 2008; Sedgley et al., 2008).

ADHESION TO TOOTH SURFACES :

Bacterial adhesion to and subsequent colonization of the surfaces of teeth and tissue are the first steps toward the formation of oral biofilms (Fig. 1). Tooth pellicle is a thin film that covers the tooth soon after the tooth is thoroughly cleansed, and it originates from salivary proteins. Oral bacteria such as Streptococci viridans can colonize the tooth surface by binding to the complex proteinaceous pellicle (Rogers et al., 2001; Kolenbrander et al., 2002). Many oral streptococci have the ability to bind to proteins such as alpha-amylase, proline-rich proteins, and proline-rich glycoproteins, and are recognized as early colonizers. Streptococcus gordonii is one of the early colonizers in dental biofilms. This species binds to acidic proline-rich proteins that account for 25-30% of the total proteins




FIG 1: A diagrammatic representation of biofilm formation on the tooth surface and the potential roles of bacterial interactions. The tooth pellicle is generally colonized by early colonizers. Co-aggregation contributes to sequential binding and colonization. Bacterial interactions include metabolic communication and genetic exchange. The development of a biofilm having a high bacterial cell density increases the concentration of signaling molecules.

in saliva. The alpha-amylase binding protein A of *S. gordonii* interacts with salivary amylase, which suggests that this interaction contributes to the attachment of *S. gordonii* to the tooth surface (Rogers et al., 2001). *S. sanguinis* is thought to be one of the first bacterial species to adhere selectively to and colonize saliva-coated teeth. This species generally appears in the human oral cavity after tooth eruption, and it becomes a normal inhabitant of the human mouth.

CO-AGGREGATION AMONG ORAL BACTERIA:

Planktonic bacterial cells that cannot directly colonize the tooth surface may bind via receptors to the cell surfaces of early colonizers that adhere to the surfaces. Co-aggregation is a specific cell-to-cell reaction that occurs between distinct bacterial cells and is one of the most important mechanisms underlying oral bacterial colonization and oral biofilm formation. Secondary colonizers bind to bacteria that are previously bound to the teeth. Sequential binding results in the appearance of a nascent surface that forms a bridge with the adjacent co-aggregating partner cells. For example, a typical periodontal pathogen, namely, *Porphyromonas gingivalis*, can bind to early colonizers. Co-adhesion between *P. gingivalis* and *S. gordonii* is mediated by two sets of adhesion–receptor pairs: the long (major) and short (minor) fimbrial subunit proteins of *P. gingivalis* that interact with streptococcal glyceraldehyde-3-phosphate dehydrogenase and Ssp surface proteins, respectively. The long (major) fimbriae of *P. gingivalis* are composed of the FimA protein, which binds to the glyceraldehyde-3-phosphate dehydrogenase present on the surface of *S. oralis* (Maeda et al., 2004). The short fimbriae of *P. gingivalis* are approximately 6.5 nm wide and 103 nm long, and are composed of the Mfa structural subunit protein. The Mfa protein engages the Ssp proteins on

the streptococcal cell surface and increases the avidity of binding to be more resistant to shear forces (Fig 2). The process of bridging between a co-aggregation of cells consisting of more than 3 bacterial species is very important, because it connects a few species that are not co-aggregation partners. *Fusobacterium nucleatum* can co-aggregate with many oral bacteria, including streptococci and obligate anaerobes. Therefore, this species is a key component of dental biofilms and serves as a coordinator that bridge the late and early colonizers (Kolenbrander et al., 2002) (Fig 3). Co-aggregation between *F. nucleatum* and other bacteria is a highly specific process involving interaction among the surface molecules of bacterial cells. The co-aggregation reactions between *F. nucleatum* and Gram-negative bacteria are mediated by lectin-carbohydrate interactions. For instance, the co-aggregation between *P. gingivalis* and *F. nucleatum* is mediated by a galactoside moiety on the surface of *P. gingivalis* and a lectin moiety on that of *F. nucleatum*, which is inhibited by lactose. Capsular polysaccharides and lipopolysaccharides of the *P. gingivalis* serotype K5 act as receptors mediating the coaggregation between oral bacteria (Rosen and Sela, 2006). In contrast, although coaggregation between *F. nucleatum* and many other Gram-positive bacteria has been observed, co-aggregation is rarely inhibited by sugars (Kolenbrander et al., 1989; Kang et al., 2005; Nagaoka et al., 2008). Thus, intergeneric co-aggregation between *F. nucleatum* and Gram-negative cells is quite different from that with Gram-positive bacteria.

Co-aggregation among oral bacteria is thought to contribute to not only bacterial colonization through physico-chemical mechanisms, but also to metabolic communication and genetic exchange, because each bacterium can easily access a neighbouring bacterial cell and its metabolites.




FIG 2: Schematic representation of differing community-relevant events that occur following the binding of *Porphyromonas gingivalis* to *Streptococcus gordonii* or to *Streptococcus cristatus*.

METABOLIC COMMUNICATION AMONG ORAL BACTERIA:

For oral bacteria, nutrients are available from saliva, gingival crevicular fluid, food containing sugars, food debris, and metabolic products of other bacteria (Fig. 4). Metabolic communications among oral bacteria may occur through the excretion of a metabolite by one organism that can be used as a nutrient by a different organism, or through the breakdown of a substrate by the extracellular enzymatic activity of one organism that creates biologically available substrates for different organisms (Kolenbrander et al., 2002). As an example, Byers et al. (1999) proposed that the hydrolysis of host glycoproteins by *S. oralis* and the subsequent utilization of released monosaccharides are important in the survival and persistence of this species and other oral bacteria. Similarly, the desialylation of immunoglobulin A1, the dominant isotype of antibody in the oral cavity, by oral Gram-positive rods may facilitate the proteolytic activities of other oral bacteria, and the concerted action may positively influence the survival of the bacteria in the oral community (Frandsen, 1994). Oral bacteria present in dental biofilms provide their metabolites as energy sources for other members. Short-chain fatty acids produced by oral bacteria are thought to be an essential carbon source for certain oral bacteria. Several studies have suggested a symbiotic association between *Streptococcus* and *Veillonella* species via lactic acid produced by the former (Kumar




FIG 3: Schematic representation of co-aggregation of primary and secondary colonizers and bridging action of *F. nucleatum*.

et al., 2005). In human studies, streptococci and veillonellae often occur in the same site of the oral cavity (Haffajee et al., 1998; Kumar et al., 2005). Moreover, interestingly, these human clinical studies suggested that these genera are associated with periodontal health. Kumar et al. (2005) stated that the parallel relationship is not surprising in view of the fact that veillonellae utilize short-chain acids such as lactates that are secreted by Gram-positive bacteria such as streptococci. From both in vitro and in vivo studies, it is probable that the growth of veillonellae present in the oral cavity depends on lactate produced by other oral bacteria. Likewise, *P. gingivalis* and *Treponema denticola* are frequently detected together in the dental biofilms of persons with periodontitis. Growth enhancement due to the mutual symbiotic relationship between *P. gingivalis* and *T. denticola* has been described in previous studies (e.g., Grenier, 1992). The growth factors produced by *P. gingivalis* and *T. denticola* were identified to be short-chain fatty acids. The growth factor produced by *T. denticola* is thought to be succinic acid, which appears to be incorporated into the lipids and phospholipids present on the cell envelope of *P. gingivalis* (Lev and Milford, 1971; Lev, 1979). Moreover, the growth of *T. denticola* is stimulated by the isobutyric acid that is produced by *P. gingivalis* (Grenier, 1992). Because these 2 species can co-aggregate, their metabolites are easily accessible to each other, since they are not diluted. Several exogenous quinones influence bacterial growth and metabolism. Vitamin K often

Table. Signaling Molecules and Functions in Oral Bacteria			
Signals	Bacteria	Functions	References
AI-2	<i>Porphyromonas gingivalis</i>	Hemin and iron uptake	Chung <i>et al.</i> (2001), James <i>et al.</i> (2006)
		Protease and hemagglutinin activities	Burgess <i>et al.</i> (2002)
		Stress gene response	Yuan <i>et al.</i> (2005)
	<i>Streptococcus mutans</i>	Biofilm formation	Yoshida <i>et al.</i> (2005)
	<i>S. gordonii</i>	Carbohydrate metabolism	McNab <i>et al.</i> (2003)
	<i>S. anginosus</i>	Biofilm formation	Petersen <i>et al.</i> (2006)
	<i>S. intermedius</i>	Virulence factors	Pecharki <i>et al.</i> (2008)
	<i>Aggregatibacter actinomycetemcomitans</i>	Biofilm formation	Shao <i>et al.</i> (2007)
	<i>Eikenella corrodens</i>	Biofilm formation	Azakami <i>et al.</i> (2006)
	<i>P. gingivalis</i> - <i>S. gordonii</i>	Biofilm formation	McNab <i>et al.</i> (2003)
CSP	<i>S. oralis</i> - <i>Actinomyces naeslundii</i>	Biofilm formation	Rickard <i>et al.</i> (2006)
	<i>S. mutans</i>	Bacteriocin production, competence	van der Ploeg (2005), Kreth <i>et al.</i> (2005b)
		Biofilm formation	Li <i>et al.</i> (2001a), Aspiras <i>et al.</i> (2004)
		Acid tolerance	Li <i>et al.</i> (2001b)
	<i>S. gordonii</i>	Biofilm formation	Loo <i>et al.</i> (2000)
	<i>S. intermedius</i>	Biofilm formation	Petersen <i>et al.</i> (2004)

sensing is considered one of the important bacterial functions. Autoinducer-2(AI-2) is one of the most well-known signaling molecules associated with quorum-sensing. It was initially identified in *Vibrio harveyi* (Bassler *et al.*, 1994) and subsequently in several bacterial species. The synthesis of AI-2 is catalyzed by LuxS, an enzyme encoded by the luxS gene. The luxS gene is highly conserved in the genome of a wide range of Gram-positive and Gram-negative bacteria. Many studies suggest that oral bacteria have a quorum-sensing system that depends on LuxS/AI-2 (Table).

Competence-stimulating peptide (CSP) mediates bacterial cell-to-cell signal communication (Table). CSP is a small soluble peptide having from 14 to 23 amino acid residues and is potentially produced by many species of oral streptococci (Li *et al.*, 2001a). CSP is implicated in bacteriocin production, virulence, and biofilm formation. Moreover, CSP enhances genetic competence, which allows for the transport of exogenous DNA into cells (Dubnau, 1991). CSP derived from *S. mutans* has been reported to induce the bacteriocin (mutacin IV) gene, and it has been shown that *S. mutans* possibly utilizes the bacteriocin to acquire the ability to transform DNA from other streptococcal species, such as *S. gordonii*, living in the same ecological niche (Kreth *et al.*, 2005b).

A new class of specifically targeted antimicrobial peptides (STAMPs) has recently been reported for use in a unique strategy (Eckert *et al.*, 2006). The STAMPs have a two-sided structure. The first is a short homing sequence of CPS that can be as unique to a bacterium as a fingerprint and ensures that the STAMPs will find their target. The second is a non-specific antibacterial peptide that is linked chemically to the homing sequence and kills the targeted bacterium on delivery. It has been suggested that STAMPs, which were designed based on the CSP of *S. mutans*, are potentially capable of eliminating *S.*

mutans from multispecies biofilms without affecting the closely related oral streptococci such as *S. gordonii* and *S. sanguinis* (Eckert *et al.*, 2006).

GENETIC EXCHANGE WITHIN COMMUNITIES:

Horizontal gene transfer by transformation, conjugation or transduction is a principal driver of bacterial evolution. The closely packed environment in biofilm communities facilitates genetic exchange among constituent cells (Nadell, 2009). The opportunistic pathogen *Pseudomonas aeruginosa*, for example, can undergo extensive genetic diversification during short-term growth in biofilm communities (Boles *et al.*, 2004). Furthermore, conjugative plasmids themselves express factors that induce their planktonic bacterial hosts to form or enter biofilm communities, which then favors the transfer of the plasmid (Ghigo, 2001). The diversity and adaptability produced by horizontal gene transfer provide a form of biological insurance that can help biofilm communities to survive in harsh environments. There are several mechanisms by which horizontal gene transfer may be operational. Mobile genetic elements like insertion sequences, transposons, integrons, bacteriophages, genomic islands, plasmids and combinations of these elements can be exchanged promiscuously between a broad spectrum of bacteria and contribute to bacterial genome plasticity. Conjugation, transformation and transduction are the other mechanism by which genetic materials are exchanged between the organisms.

CONCLUSION

The oral biofilm is sophisticated social networking, based initially on very specific

recognition of surface characteristics, which provides the discrimination necessary for the formation of metabolically compatible, physiologically integrated communities. Community development is controlled by programmed patterns of gene expression and multilevel regulation of protein expression and activity. Organisms within these communities continually monitor the host environment and the nature and intentions of other organisms that may seek to participate in community affairs. Interspecies communication, thus maintain a dynamic and stable environment. Once a degree of stability or maturity is reached, organisms can begin the process of genetic exchange and the production of genetically diverse daughter cells, some of which will exhibit increased fitness. This phenomenon makes oral biofilm to survive throughout life since its inception.

REFERENCES

1. Bassler BL, Wright M, Silverman MR (1994). Multiple signalling systems controlling expression of luminescence in *Vibrio harveyi*: sequence and function of genes encoding a second sensory pathway. *Mol Microbiol* 13:273-286.
2. Grenier D (1992). Nutritional interactions between two suspected periodontopathogens, *Treponema denticola* and *Porphyromonas gingivalis*. *Infect Immun* 60:5298-5301.
3. Boles BR, Thoendel M, Singh PK. Self-generated diversity produces "insurance effects" in biofilm communities. *Proc Natl Acad Sci U S A* 2004;101: 16630–16635.
4. Byers HL, Tarelli E, Homer KA, Beighton D (1999). Sequential deglycosylation and utilization of the N-linked, complex-type glycans of human alpha1-acid glycoprotein mediates growth of *Streptococcus oralis*. *Glycobiology* 9:469-479.
5. Chalmers NI, Palmer RJ, Cisar JO, Kolenbrander PE (2008). Characterization of a *Streptococcus* sp.-*Veillonella* sp. community micromanipulated from dental plaque. *J Bacteriol* 190:8145-8154.
6. Dashper SG, Reynolds EC (2000). Effects of organic acid anions on growth, glycolysis, and intracellular pH of oral streptococci. *J Dent Res* 79:90-96.
7. Dubnau D (1991). Genetic competence in *Bacillus subtilis*. *Microbiol Rev* 55:395-424.
8. Eckert R, He J, Yarbrough DK, Qi F, Anderson MH, Shi W (2006). Targeted killing of *Streptococcus mutans* by a pheromone-guided "smart" antimicrobial peptide. *Antimicrob Agents Chemother* 50:3651-3657.
9. Ghigo JM. Natural conjugative plasmids induce bacterial biofilm development. *Nature* 2001;412: 442–445.
10. Haffajee AD, Cugini MA, Tanner A, Pollack RP, Smith C, Kent RL Jr, et al. (1998). Subgingival microbiota in healthy, well-maintained elder and periodontitis subjects. *J Clin Periodontol* 25:346-353.
11. Isawa K, Hojo K, Yoda N, Kamiyama T, Makino S, Saito M, et al. (2002). Isolation and identification of a new bifidogenic growth stimulator produced by *Propionibacterium freudenreichii* ET-3. *Biosci Biotechnol Biochem* 66:679-681.
12. Kang MS, Na HS, Oh JS (2005). Coaggregation ability of *Weissella cibaria* isolates with *Fusobacterium nucleatum* and their adhesiveness to epithelial cells. *FEMS Microbiol Lett* 253:323-329
13. Kolenbrander PE, Andersen RN, Moore LV (1989). Coaggregation of *Fusobacterium nucleatum*, *Selenomonas flueggei*, *Selenomonas infelix*, *Selenomonas noxia*, and *Selenomonas sputigena* with strains from 11 genera of oral bacteria. *Infect Immun* 57:3194-3203.
14. Kolenbrander PE, Andersen RN, Blehert DS, Eglund PG, Foster JS, Palmer RJ Jr (2002). Communication among oral bacteria. *Microbiol Mol Biol Rev* 66:486-505.
15. Kreth J, Merritt J, Shi W, Qi F (2005a). Competition and coexistence between *Streptococcus mutans* and *Streptococcus sanguinis* in the dental biofilm. *J Bacteriol* 187:7193-7203.
16. Kumar PS, Griffen AL, Moeschberger ML, Leys EJ (2005). Identification of candidate periodontal pathogens and beneficial species by quantitative 16S clonal analysis. *J Clin Microbiol* 43:3944-3955.
17. Lev M, Milford AF (1971). Vitamin K stimulation of sphingolipid synthesis. *Biochem Biophys Res Commun* 45:358-362.
18. Lev M (1979). Sphingolipid biosynthesis and vitamin K metabolism in *Bacteroides melaninogenicus*. *Am J Clin Nutr* 32:179-186.
19. Li YH, Hanna MN, Svensäter G, Ellen RP, Cvitkovitch DG (2001b). Cell density modulates acid adaptation in *Streptococcus mutans*: implications for survival in biofilms. *J Bacteriol* 183:6875-6884.
20. Marcotte H, Lavoie MC (1998). Oral microbial ecology and the role of salivary immunoglobulin A. *Microbiol Mol Biol Rev* 62:71-109.
21. Maeda K, Nagata H, Yamamoto Y, Tanaka M, Tanaka J, Minamino N, et al. (2004). Glyceraldehyde-3-phosphate dehydrogenase of *Streptococcus oralis* functions as a coadhesin for *Porphyromonas gingivalis* major fimbriae. *Infect Immun* 72:1341-1348.
22. Nadell CD, Xavier JB, Foster KR. The sociobiology of biofilms. *FEMS Microbiol Rev* 2009;33: 206–224.
23. Nagaoka S, Hojo K, Murata S, Mori T, Ohshima T, Maeda N (2008). Interactions between salivary *Bifidobacterium adolescentis* and other oral bacteria:

in vitro coaggregation and coadhesion assays. FEMS Microbiol Lett 281:183-189.

23. Pangsomboon K, Kaewnopparat S, Pitakpornpreecha T, Srichana T (2006). Antibacterial activity of a bacteriocin from *Lactobacillus paracasei* HL32 against *Porphyromonas gingivalis*. Arch Oral Biol 51:784-793.

24. Pangsomboon K, Bansal S, Martin GP, Sutinanalert P, Kaewnopparat S, Srichana T (2009). Further characterization of a bacteriocin produced by *Lactobacillus paracasei* HL32. J Appl Microbiol 106:1928-1940.

25. Rogers JD, Palmer RJ, Kolenbrander PE, Scannapieco FA (2001). Role of *Streptococcus gordonii* amylase-binding protein A in adhesion to hydroxyapatite, starch metabolism, and biofilm formation. Infect Immun 69:7046-7056.

26. Rosen G, Sela MN (2006). Coaggregation of *Porphyromonas gingivalis* and *Fusobacterium nucleatum* PK 1594 is mediated by capsular polysaccharide and lipopolysaccharide. FEMS Microbiol Lett 256:304-310.

27. Sedgley CM, Lee EH, Martin MJ, Flannagan SE (2008). Antibiotic resistance gene transfer between *Streptococcus gordonii* and *Enterococcus faecalis* in root canals of teeth ex vivo. J Endod 34:570-574.